FORM 4 TERM 3 APRIL 2022 CHEMISTRY PAPER 3

You are provided with

- Anhydrous sodium carbonate solid x.
- Distilled water.
- 0.2m Hydrochloric acid solution A.

You are required to determine molar heat of solution of solid x.

PROCEDURE I

- i. Place 50.0ml of water in 250ml plastic beaker.
- ii. Note the temperature of the water and record it in the table I below.
- iii. Add all the solid X provided to the water in the beaker, stir gently with the thermometer and record the final temperature of the solution in the table I below. Keep the resulting solution for procedure 2.

TABLE I

Final temperature (⁰ C)	
Initial temperature (⁰ C)	
Change in temperature (⁰ C)	

(2 mks)

(a) What is the enthalpy change for the reaction? (Assume the density of solution is 1g/cm³, and specific heat capacity is 4.2 Jg⁻¹ K⁻¹). (2 mks)

PROCEDURE II

Transfer the contents of the beaker into 250ml volumetric flask. Rinse both the beaker and the thermometer with distilled water and ass this water into the solution in the volumetric flask. Add more water to make up to the mark. Label this solution as solution X. fill the burette with solution A. Using a pipette place 25.0ml of solution X into a conical flask. Add 3 drops of methyl orange indicator and titrate with solution A. record your readings in table II below. Repeat the titration two more times and complete the table.

TABLE II

Experiment		
Final burette reading (cm ³)		
Initial burette reading (cm ³)		
Volume of solution A used (cm ³)		

(3 mks)

(b) Calculate average volume of solution A used. (1 mk)

(c) the number of moles of solution A used.	(1 mk)
(d) The number of moles of solution X that reacted with the number of moles of s (c) above.	olution A in (1 mk)
(e) The number of moles of solid X used in procedure I.	(1 mk)
(f) Molar heat of solution of anhydrous sodium carbonate.	(2 mks)

- 2. You are provided with:
- A solution of sodium hydroxide labeled B.
- A solution of sulphuric(vi)acid labeled C.

You are required to determine the concentration of the alkali using the following procedure.

PROCEDURE:

- (i) Place 40cm³ of sodium hydroxide solution into a 250 ml plastic beaker.
- (ii) Measure 60cm³ of sulphuric (vi) acid solution.
- (iii) Determine the temperature of sodium hydroxide solution at half a minute intervals for two minutes and record it in the table below.
- (iv) At $2\frac{1}{2}$ minutes, place the 60cm^3 of solution C into the plastic beaker while stirring and resume taking the temperature in the 3^{rd} minute.
- (v) Complete the table below.

Time in minutes	0	1/2	1	1 ½	2	2 ½	3	3 ½	4

Temperature in ⁰ C							
		1	Ī		1	T	
Time in minutes	4 ½	5	5 ½	6	6 ½	7	
Temperature in ⁰ C							
							(3 mks)
(a) Plot a graph of tem	perature aga	ninst time.					(3 mks)
(b) From the graph, determine the highest temperature change.							(1 mk)
(c) Determine the heat evolved in this experiment (Density of solution = 1 g/cm³ specific heat capacity of solution = 4.2 Jg-¹ K-¹) (2 mks) (d) Given that the molar heat of neutralization is 56KJ/mole, determine the number of mole of sodium hydroxide used in the neutralization reaction above. (2 mks)							
(e) Determine the mola	nrity of sodi	um hydro	xide.				(2 mks)

3. You are provided with solid K. carry out the following tests and write your observations and inferences in the spaces provided.

	TEST	OBSERVATION	INFERENCE
(a)	Place a spatula full of sample K in a		
	clean dry test tube. Heat gently and		
	then strongly.	(1 mk)	(1 mk)
(b)	Put the remaining solid K in a boiling		
	tube. Add about 8cm ³ of distilled		
	water. Shake well and divide the		
	solution into 3 portions.		
(i)	To the first portion add 3 drops of		
	sodium hydroxide solution and then		
	excess.	(1 mk)	(1 mk)
(ii)	To the second portion add 3 drops of		
	ammonia solution and then excess.		
		(1 mk)	(1 mk)
(iii)	To the third portion add 3 drops of		

	Barium nitrate followed by 3 drops of		
	nitric acid.	(1 mk)	(1 mk)
(c)	You are provided with solid P. carry out	the tests below and record y	our observations and
	inferences.		
(i)	Place half spatula of solid P in a non-		
	luminous flame of a Bunsen burner.		
		(1 mk)	(1 mk)
(ii)	Dissolve the remaining solid in water		
	and divide into two portions		
(a)	Add 3 drops of universal indicator to		
	the 1^{st} portion and determine the PH of the solution.	(1 mk)	(1 mk)
(b)	To the 2 nd portion add a little sodium		
	hydrogen carbonate		

(1 mk)	(1 mk)